

GEOMETRY OF QUANTUM HOMOGENEOUS SUPERVECTOR BUNDLES AND REPRESENTATION THEORY OF QUANTUM GENERAL LINEAR SUPERGROUP

R. B. ZHANG

Department of Pure Mathematics
University of Adelaide
North Terrace, Adelaide
S.A. 5001, Australia

The quantum general linear supergroup $GL_q(m|n)$ is defined and its structure is studied systematically. Quantum homogeneous supervector bundles are introduced following Connes' theory, and applied to develop the representation theory of $GL_q(m|n)$. Quantum Frobenius reciprocity is proven, and a Borel - Weil theorem is established for the covariant and contravariant tensor irreps.

Quantum supergroup $GL_q(m|n)$

We will work on the complex field $\mathbf{C}$. For convenience, the Lie superalgebra $gl(m|n)$ will be denoted by $\mathbf{g}$, and its quantized universal enveloping algebra by $U_q(\mathbf{g})$. Here $U_q(\mathbf{g})$ is of Jimbo type with q being specialized to a generic complex parameter. We will largely follow the notations of [1] and [2]. Now the generators of $U_q(\mathbf{g})$ are $K_a, K_a^{-1}, E_{b-b+1}, E_{b+1,b}, a \in \mathbf{I}, b \in \mathbf{I}'$, where $\mathbf{I} = \{1, 2, \dots, m+n\}$, $\mathbf{I}' = \mathbf{I} \setminus \{m+n\}$, and the defining relations are given in [1]. Corresponding to any subset Θ of $\mathbf{I}'$, we introduce $U_q(\mathbf{k})$ which is generated by the elements of $\mathcal{S}_k = \{K_a^{\pm 1}, E_{c-c+1}, E_{c+1,c}, a \in \mathbf{I}, c \in \Theta\}$, and $U_q(\mathbf{p}_{\pm})$ respectively generated by those of $\mathcal{S}_k \cup \{E_{c-c+1}, c \in \mathbf{I}' \setminus \Theta\}$ and $\mathcal{S}_k \cup \{E_{c+1,c}, c \in \mathbf{I}' \setminus \Theta\}$. Clearly, $U_q(\mathbf{p}_{\pm})$ are $\mathbf{Z}_2$ -graded Hopf subalgebras of $U_q(\mathbf{g})$, which will be called parabolic. Also, $U_q(\mathbf{k})$ is a $\mathbf{Z}_2$ -graded Hopf subalgebra of $U_q(\mathbf{p}_{\pm})$.

The representation theory of $U_q(\mathbf{g})$ has been systematically developed [1]. Let π be the contravariant vector representation of $U_q(\mathbf{g})$ afforded by the irreducible module E in the standard basis $\{v_a\}$ such that $E_{a-a\pm 1}v_b = \delta_{b-a\pm 1}v_a$, $K_av_b = q_a^{\delta_{ab}}v_b$. Although finite dimensional representations of $U_q(\mathbf{g})$ are not completely reducible in general, repeated tensor products of π can be reduced into direct sums of contravariant tensor irreps. Let $\Lambda^{(1)}$ be the set of the highest weights of all such irreps. Let $\overline{E}$ be the dual module of E , and $\overline{\pi}$ the covariant vector representation (dual to π). Repeated tensor products of $\overline{\pi}$ are again completely reducible. We denote by $\Lambda^{(2)}$ the highest weights of all the covariant tensor irreps. (The trivial representation is tensorial by convention.) $\Lambda^{(1)}$ and $\Lambda^{(2)}$ can be explicitly characterized [3]. In particular, it is known that $\Lambda^{(1)} \cap \Lambda^{(2)} = \{0\}$. Also, if $\lambda \in \Lambda^{(1)}$, we denote by $\bar{\lambda}$ the lowest weight of the irrep with highest weight λ , and set $\lambda^\dagger = -\bar{\lambda}$. Then $\Lambda^{(2)} = \{\lambda^\dagger | \lambda \in \Lambda^{(1)}\}$.

It follows from standard Hopf algebra theory that $(U_q(\mathbf{g}))^0$ is a Hopf superalgebra with a structure dualizing that of $U_q(\mathbf{g})$. Consider $T_q \subset (U_q(\mathbf{g}))^0$ generated by the matrix elements of π : $t_{ab} \in (U_q(\mathbf{g}))^0, a, b \in \mathbf{I}$. It is a $\mathbf{Z}_2$ -graded bi-subalgebra of $(U_q(\mathbf{g}))^0$ with the generators satisfying an 'RTT' relation [2], but does not admit an antipode. Our earlier discussions on representations of $U_q(\mathbf{g})$ imply that the matrix elements of the contravariant tensor irreps form a Peter-Weyl basis for T_q . Similarly, the matrix elements $\bar{t}_{ab} \in (U_q(\mathbf{g}))^0$ of $\overline{\pi}$ also generate a $\mathbf{Z}_2$ -graded bi-subalgebra $\overline{T}_q$ of $(U_q(\mathbf{g}))^0$, for which the matrix elements of the covariant tensor irreps form a Peter-

Weyl basis.

We define the algebra of functions G_q on $GL_q(m|n)$ to be the subalgebra of $(U_q(\mathfrak{g}))^0$ generated by $\{t_{ab}, \bar{t}_{ab} | a, b \in \mathbf{I}\}$. It inherits a bi - superalgebra structure from T_q and $\bar{T}_q$, and also admits an antipode. Thus G_q is a Hopf superalgebra. By considering the universal R - matrix of $U_q(\mathfrak{g})$ we can easily show that the following relation is satisfied in G_q

$$R_{12}^{\bar{\pi}\pi} \bar{t}_1 t_2 = t_2 \bar{t}_1 R_{12}^{\bar{\pi}\pi}, \quad (1)$$

where $R^{\bar{\pi}\pi} = q^{-\sum_{a \in \mathbf{I}} e_{aa} \otimes e_{aa} (-1)^{[a]}} - (q - q^{-1}) \sum_{a < b} e_{ba} \otimes e_{ba} (-1)^{[a] + [b] + [a][b]}$. An immediate consequence of (1) is the factorization $G_q = T_q \bar{T}_q$. It can also be shown [2] that G_q separates points of $U_q(\mathfrak{g})$, that is, for any nonvanishing $x \in U_q(\mathfrak{g})$, there exists $f \in G_q$ such that $f(x) \neq 0$.

Induced representations and quantum super-vector bundles

Let us introduce two left actions L and R of $U_q(\mathfrak{g})$ on G_q . For any $x \in U_q(\mathfrak{g})$ and $f \in G_q$, we define $R_x(f) = \sum_{(f)} (-1)^{[x]([f] + [x])} f_{(1)} \langle f_{(2)}, x \rangle$ and $L_x(f) = \sum_{(f)} \langle f_{(1)}, S^{-1}(x) \rangle f_{(2)}$, where Sweedler's sigma notation is employed. We *trivially* extend them to $V \otimes_{\mathbf{C}} G_q$, and still denote the resultant actions by the same notations. Clearly, L and R graded - commute. Now define

$$\mathcal{A}_q := \{f \in G_q | R_y(f) = \epsilon(y)f, \forall y \in U_q(\mathbf{k})\}, \quad (2)$$

which forms a subalgebra of G_q . It can be regarded as the superalgebra of functions on a quantum homogeneous superspace. Given any finite dimensional $U_q(\mathbf{k})$ - module V , we define

$$\mathcal{E}(V) := \{\zeta \in V \otimes_{\mathbf{C}} G_q | R_y(\zeta) = (S(y) \otimes id_{G_q})\zeta, \forall y \in U_q(\mathbf{k})\}. \quad (3)$$

Then it immediately follows from the commutativity of R and L that $\mathcal{E}(V)$ furnishes a left $U_q(\mathfrak{g})$ - module under L . We will call $\mathcal{E}(V)$ an induced $U_q(\mathfrak{g})$ - module. There exists the following quantum Frobenius reciprocity: *if W is a quotient $U_q(\mathfrak{g})$ - module of $\bigoplus_{k,l} \{E^{\otimes k} \otimes \bar{E}^{\otimes l}\}$, then there is the canonical isomorphism*

$$Hom_{U_q(\mathfrak{g})}(W, \mathcal{E}(V)) \cong Hom_{U_q(\mathbf{k})}(W, V). \quad (4)$$

To understand the underlying geometry of induced representations, observe that $\mathcal{E}(V)$ forms a two - sided $\mathcal{A}_q$ - module under the multiplication of G_q . If W is a $U_q(\mathfrak{g})$ - module, then $\mathcal{E}(W)$ is free. It follows that if V is a $U_q(\mathfrak{k})$ - module which is contained as a direct summand in a $U_q(\mathfrak{g})$ - module, then $\mathcal{E}(V)$ is project, and in this case, we may regard $\mathcal{E}(V)$ as the space of sections of a quantum supervector bundle [4] associated with $\mathcal{A}_q$. When $m \notin \Theta$, every finite dimensional $U_q(\mathfrak{k})$ - module V has this property. Also, for general Θ , $\mathcal{E}(V)$ is projective if it yields non - zero $\mathcal{O}(\mu)$ or $\overline{\mathcal{O}}(\mu)$ (defined below).

Let V be a finite dimensional irreducible module over $U_q(\mathfrak{p}) = U_q(\mathfrak{p}_\pm)$ with the $U_q(\mathfrak{k})$ highest weight μ and lowest weight $\tilde{\mu}$. Define the subspaces $\mathcal{O}(\mu)$ and $\overline{\mathcal{O}}(\mu)$ of $\mathcal{E}(V)$ by

$$\begin{aligned}\overline{\mathcal{O}}(\mu) &:= \{\zeta \in \mathcal{E}(V) \cap (V \otimes T_q) \mid R_y(\zeta) = (S(y) \otimes id_{G_q})\zeta, \forall y \in U_q(\mathfrak{p})\}; \\ \mathcal{O}(\mu) &:= \{\zeta \in \mathcal{E}(V) \cap (V \otimes \overline{T}_q) \mid R_y(\zeta) = (S(y) \otimes id_{G_q})\zeta, \forall y \in U_q(\mathfrak{p})\}.\end{aligned}\quad (5)$$

Then we have the following quantum Borel - Weil theorem for the covariant and contravariant tensor irreps [2]: *As $U_q(\mathfrak{g})$ - modules,*

$$\begin{aligned}\mathcal{O}(\mu) &\cong \begin{cases} W((- \tilde{\mu})^\dagger), & \text{if } \tilde{\mu} \in -\Lambda^{(2)}, \quad U_q(\mathfrak{p}) = U_q(\mathfrak{p}_+), \\ W(\mu), & \text{if } \mu \in \Lambda^{(1)}, \quad U_q(\mathfrak{p}) = U_q(\mathfrak{p}_-), \\ \{0\}, & \text{otherwise.} \end{cases} \\ \overline{\mathcal{O}}(\mu) &\cong \begin{cases} W((- \tilde{\mu})^\dagger), & \text{if } \tilde{\mu} \in -\Lambda^{(1)}, \quad U_q(\mathfrak{p}) = U_q(\mathfrak{p}_+), \\ W(\mu), & \text{if } \mu \in \Lambda^{(2)}, \quad U_q(\mathfrak{p}) = U_q(\mathfrak{p}_-), \\ \{0\}, & \text{otherwise,} \end{cases}\end{aligned}\quad (6)$$

where the notation $W(\lambda)$ signifies the irreducible $U_q(\mathfrak{g})$ - module with highest weight λ .

References

- [1] R. B. Zhang, J. Math. Phys., **34**(1993) 1236.
- [2] R. B. Zhang, Commun. Math. Phys., in press (q-alg/9611024).
- [3] M. D. Gould and R. B. Zhang, J. Math. Phys., **31** (1990) 2552.
- [4] A. Connes, *Noncommutative geometry*, Academic Press (1994).